



Name:


2. Cinnamaldehyde can be reduced using sodium borohydride, NaBH<sub>4</sub>.

(i) State which functional group reacts with the sodium borohydride.

.....

[1]


(ii) Complete and balance the equation for this reaction.



[1]

[Total 2 marks]

3. The reducing agent, NaBH<sub>4</sub>, is used widely in organic chemistry. One example is for the reduction of diphenylethanedione, C<sub>14</sub>H<sub>10</sub>O<sub>2</sub>, shown below.



**diphenylethanedione**

(i) Draw a displayed formula to show the structure of the organic product that would be formed by reducing diphenylethanedione with excess NaBH<sub>4</sub>.

[1]

Name:

- (ii) Complete and balance the equation for this reaction, using [H] to represent the reducing agent.




[1]

[Total 2 marks]

4. Chloral hydrate is broken down in the body after several hours. One reaction is oxidation to trichloroethanoic acid.

Complete the equation for this reaction below.



[Total 1 mark]

5. But-2-enal,  $\text{CH}_3\text{CH}=\text{CHCHO}$ , is a pale yellow, flammable liquid with an irritating odour.

- (a) (i) Describe a simple chemical test that would show that but-2-enal is an aldehyde.

.....  
.....  
.....

[2]

- (ii) Explain why this test gives a different result with aldehydes than it does with ketones.

.....  
.....

[1]

Name:

(b) But-2-enal also reacts with sodium borohydride,  $\text{NaBH}_4$ .

(i) Identify the organic compound formed in this reaction.

.....

[1]

(ii) State the type of chemical reaction occurring.

.....

[1]

(c) Precautions must be taken to prevent but-2-enal catching fire.


Construct a balanced equation for the complete combustion of but-2-enal,  
 $\text{C}_4\text{H}_6\text{O}$ .

[1]


[Total 6 marks]

Name:


6. An unknown colourless liquid with molecular formula  $C_4H_8O$  was thought to be one of butanal, but-3-en-1-ol, or butanone.



**butanal**



**but-3-en-1-ol**



**butanone**

- (a) State a simple chemical test that would positively identify:

- (i) butanal **only**;

reagent .....

observation .....

organic product .....

[3]

- (ii) but-3-en-1-ol **only**.

reagent .....

observation .....

type of reaction .....

[3]

Name:

- (b) Butanal and butanone both react with 2,4-dinitrophenylhydrazine to produce mixtures containing orange precipitates.

Outline how the mixtures containing these orange precipitates can be used to distinguish between butanal and butanone.

.....  
.....  
.....  
.....  
.....  
.....

[3]

[Total 9 marks]

7. Hydroxyethanal,  $\text{HOCH}_2\text{CHO}$ , is sometimes referred to as the 'first sugar' as it is the simplest possible molecule that contains both an aldehyde group and an alcohol group.

A biochemist investigated some redox reactions of hydroxyethanal and found that several different products were produced.

- (a) The biochemist reacted hydroxyethanal with Tollens' reagent.

- (i) State what the biochemist would see when hydroxyethanal reacts with Tollens' reagent.

.....

[1]

- (ii) Write the structural formula of the organic product formed when hydroxyethanal reacts with Tollens' reagent.

[1]

Name:

- (b) The biochemist also reacted hydroxyethanal with acidified dichromate by heating under reflux.

Write an equation for this oxidation.

Use **[O]** to represent the oxidising agent.

[2]

- (c) The biochemist then reduced hydroxyethanal using aqueous  $\text{NaBH}_4$ .

- (i) Write the structural formula of the organic product.

.....

[1]

- (ii) Outline the mechanism for this reduction.

Use curly arrows and show any relevant dipoles.

[4]

[Total 9 marks]